

A construction of a λ -Poisson generic sequence

Gabriel Sac Himelfarb

1 Abstract

Years ago Zeev Rudnick defined the λ -Poisson generic sequences as the infinite sequences of symbols in a finite alphabet where the number of occurrences of long words in the initial segments follow the Poisson distribution with parameter λ . Benjamin Weiss and Yuval Peres [5] proved that almost all sequences with respect to Lebesgue measure are Poisson generic (see also [1]). Despite this result, no explicit example has yet been given.

In this talk I will present a construction of an explicit λ -Poisson generic sequence over an alphabet of at least three symbols, for any fixed positive real number λ . This is joint work with Verónica Becher. Since λ -Poisson genericity implies Borel normality, the constructed sequence is Borel normal. The same construction provides explicit instances of Borel normal sequences that are not λ -Poisson generic.

Given $x \in \Omega^{\mathbb{N}}$, and a positive real number λ , $i \in \mathbb{N}_0$ and $k \in \mathbb{N}$ we write $Z_{i,k}^\lambda(x)$ for the proportion of words of length k that occur exactly i times in the prefix of x of length $\lfloor \lambda b^k \rfloor$.

Definition. Let λ be a positive real number. A sequence $x \in \Omega^{\mathbb{N}}$ is λ -Poisson generic if for every $i \in \mathbb{N}_0$,

$$\lim_{k \rightarrow \infty} Z_{i,k}^\lambda(x) = e^{-\lambda} \frac{\lambda^i}{i!}.$$

A sequence is Poisson generic if it is λ -Poisson generic for all positive real numbers λ .

The main result I will discuss is the following:

Theorem 1 ([Becher and S.H. [3, Theorem 1]]). *Let λ be a positive real number and Ω a b -symbol alphabet, $b \geq 3$. Let $(p_i)_{i \in \mathbb{N}_0}$ be a sequence of non-negative real numbers such that $\sum_{i \geq 0} p_i = 1$ and $\sum_{i \geq 0} i p_i = \lambda$. Then there is a construction of an infinite sequence x over alphabet Ω , which satisfies for every $i \in \mathbb{N}_0$,*

$$\lim_{k \rightarrow \infty} Z_{i,k}^\lambda(x) = p_i.$$

The construction in Theorem 1 consists in concatenating segments of any infinite de Bruijn sequence [2, Theorem 1], which is a sequence that satisfies that each initial segment of length b^k is a cyclic de Bruijn sequence of order k . Our construction works by selecting segments of this given sequence and repeating them as many times as determined by the probabilities p_i , for every $i \in \mathbb{N}_0$.

Weiss showed that 1-Poisson genericity implies Borel normality and that the two notions do not coincide¹, witnessed by the Champernowne sequence [4]. The next result is a normality

¹Benjamin Weiss. Random-like behavior in deterministic systems, 16 June 2010. Institute for Advanced Study Princeton University USA. The conference is available at https://www.youtube.com/watch?v=8AB7591De68&ab_channel=InstituteForAdvancedStudy

criterion that generalizes this fact. In contrast to Theorem 1, this result does not require the alphabet size b to be greater than 2.

Theorem 2 ([Becher and S.H. [3, Theorem 2]). *Let Ω be a b -symbol alphabet, $b \geq 2$, and let $x \in \Omega^{\mathbb{N}}$. We fix a positive real number λ and define for every $i \in \mathbb{N}_0$ the numbers $p_i = \liminf_{k \rightarrow \infty} Z_{i,k}^\lambda(x)$. If the numbers p_i satisfy $\sum_{i \geq 0} ip_i = \lambda$ then x is normal to base b .*

As a consequence of Theorem 2,

Corollary 1. *Every λ -Poisson generic sequence is Borel normal, but the two notions do not coincide. The construction in Theorem 1 yields infinitely many Borel normal sequences which are not λ -Poisson generic.*

References

- [1] Nicolás Alvarez, Verónica Becher, and Martín Mereb. Poisson generic sequences. *Submitted*, February 4, 2022. Preprint <https://arxiv.org/abs/2202.01632>.
- [2] Verónica Becher and Pablo Ariel Heiber. On extending de Bruijn sequences. *Information Processing Letters*, 111:930–932, 2011.
- [3] Verónica Becher and Gabriel Sac Himelfarb. A construction of a λ - poisson generic sequence. *Submitted*, May 9, 2022. Preprint <https://arxiv.org/abs/2205.03981>.
- [4] David Champernowne. The construction of decimals normal in the scale of ten. *Journal of London Mathematical Society*, s1-8(4):254–260, 1933.
- [5] Benjamin Weiss. Poisson generic points. Jean-Morlet Chair 2020 - Conference: Diophantine Problems, Determinism and Randomness, Centre International de Rencontres Mathématiques, November 23 to 29, 2020. Audio-visual resource: doi:10.24350/CIRM.V.19690103.